Promotion of health education on AIDS prevention

In DECEMBER of 2019

A. Regular Inmates

1. New inmates:

Playing sanitation education promotional video on AIDS during investigation of new inmates to provide all inmates an initial knowledge on AIDS. (2 times , for a total of67 person-time)

2. In-prison inmates:

Invite professional lecturers every month in a regular cyclical format to sponsor AIDS prevention class in each factory and ward to instill the correct concept and knowledge about AIDS and to teach inmates on things to watch out for and direction of future life after being released.

(1time , for a total of 96 person-time)




3. Promotion of sanitation education to regular inmates:

education speech

Video conference while in clinic with materials of sanitation education promotion, or distributing sanitation education materials on AIDS which are also distributed to every unit in order to increase the knowledge and awareness in the inmates about the disease.

(6 times , for a total of 121person-time)



B. The Infected

1. Newly infected :

The public health nurse of Yunlin Public Health Bureau or disease control individual controller provide the sanitation education to get to know the disease, adjust the mood of the inmates, and provided information to the contacted to prevent the spread of the disease.

(no new individual inmate case)

2. In-prison infected inmates :

The volunteers from Taiwan Lourdes Association provide monthly group counseling with story reading, singing hymns, drawing, and group activities in a sustained fashion in caring for the body and soul of the infected inmates. They also arranged religious classes to give support from different approach.



(1time, for a total of 3 person-time)

Professional individual sanitation education ducation 3. About to be released infected inmates :

The public health nurse of Yunlin Public Health Bureau or disease control individual controller conduct individual interviews to help with after-release medical care and settlement so that the infected can successfully maintain contact with social support group and reduce the chance of using narcotics or infecting others.

