Promotion of health education on AIDS prevention in April of 2021

A. Regular Inmates

1. New inmates:

Playing sanitation education promotional video on AIDS during investigation of new inmates to provide all inmates an initial knowledge on AIDS.

(1 times for a total of 9 person-time)

2. In-prison inmates:

Invite professional lecturers every month in a regular cyclical format to sponsor AIDS prevention class in each factory and ward to instill the correct concept and knowledge about AIDS and to teach inmates on things to watch out for and direction of future life after being released.

(1 time for a total of 77 person-time)



Brilliant professional sanitation education speech



attentively listening

3. Promotion of sanitation education to regular inmates:
Video conference while in clinic with materials of sanitation education
promotion, or distributing sanitation education materials on AIDS which are also
distributed to every unit in order to increase the knowledge and awareness in
the inmates about the disease.

(5times for a total of 107 Person-time)



VCD



B. The Infected

1. Newly infected:

The public health nurse of Yunlin Public Health Bureau or disease control individual controller provide the sanitation education to get to know the disease, adjust the mood of the inmates, and provided information to the contacted to prevent the spread of the disease.

(0 time for a total of 0 person-time)

2. In-prison infected inmates:

The volunteers from Taiwan Lourdes Association provide monthly group counseling with story reading, singing hymns, drawing, and group activities in a sustained fashion in caring for the body and soul of the infected inmates. They also arranged religious classes to give support from different approach.

(1 time for a total of 2 person-time)



3. About to be released infected inmates:

The public health nurse of Yunlin Public Health Bureau or Social worker conduct individual interviews to help with after-release medical care and settlement so that the infected can successfully maintain contact with social support group and reduce the chance of using narcotics or infecting others.

(1 time for a total of 1 person-time)

